The development of new soybean strain with *ti* and *cgy₁* recessive allele

Sang Woo Choi • Jun Hyun Park • Jong Il Chung

Received: 2 August 2018 / Revised: 17 September 2018 / Revised: 23 October 2018 / Accepted: 23 October 2018
© Korean Society for Plant Biotechnology

Abstract Soybean [*Glycine max* (L.) Merr.] seed is an important dietary source of protein, oil, carbohydrate, isoflavone and other various nutrients for humans and animals. However, there are anti-nutritional factors in the raw mature soybeans. Kunitz trypsin inhibitor (KTI) protein and stachyose are the main anti-nutritional factors in soybean seed. The α' -subunit of β -conglycinin protein exhibit poor nutritional and food processing properties. The genetic removal of the KTI and α' -subunit proteins will improve the nutritional value of the soybean seed. The objective of this research was to develop a new soybean strain with KTI and α' -subunit protein free (*titicgy₁cgy₁* genotype) and proper agronomic traits. A breeding population was developed from the cross of the BL-1 and 15G1 parents. A total of 168 F₂ seeds from the cross of the BL-1 and 15G1 parents were obtained. The segregation ratios of 9: 3: 3: 1 (104 *Ti_Cgy₁*_: 30 *Ti_cgy₁cgy₁*: 21 *cgy₁cgy₁Ti*_: 13 *titicgy₁cgy₁*) between the *Ti* and *Cgy₁* genes in the F₂ seeds were observed ($\chi^2=5.12$, $P=0.5-0.10$). Two F₄ plant strains with proper agronomical traits and *titicgy₁cgy₁* genotype (free of both KTI and α' -subunit protein) were selected and harvested. 2 strains (S1 and S2) had yellow seed coats and hilum. The plant height of the S1 strain was 65 centimeters. The 100-seed weight was 29.2 g. The plant height of the S2 strain was 66 centimeters and 100-seed weight was 26.2 g. The two strains selected in this research will be used to improve the new cultivar that will be free of the KTI and α' -subunit proteins.

Keywords Kunitz trypsin inhibitor (KTI), α' -subunit, breeding lines, soybean

Introduction

Soybean [*Glycine max* (L.) Merr.] seeds contain protein, carbohydrate, oil, vitamins, minerals, and many functional components. Also, there are antinutritional factors and several allergenic proteins in the raw mature soybean. Kunitz trypsin inhibitor (KTI) protein of mature soybean seed is a main antinutritional factor in soybean seed. KTI protein is a small and non-glycosylated protein containing 181 amino acid residues with 21.5 kDa. KTI protein was first isolated and crystallized from soybean seeds by Kunitz (1945). Kunitz trypsin inhibitor protein strongly inhibits trypsin, thus reducing food intake by diminishing digestion and absorption. Five electrophoretic forms of KTI have been discovered. The genetic control of four forms, *Ti^a*, *Ti^b*, *Ti^c*, and *Ti^d*, has been reported as a codominant multiple allelic series at a single locus (Sing et al. 1969; Hymowitz and Hadley 1972; Orf and Hymowitz 1979). Orf and Hymowitz (1979) found that the fifth form does not exhibit a soybean trypsin inhibitor-A2 band and is inherited as a recessive allele designated *ti*. Orf and Hymowitz (1979) identified two soybean accessions (PI157440 and PI196168) which lacks KTI protein from USDA germplasm collection. The *Ti* locus has been located on linkage group 9 in the classical linkage map of soybean (Hildebrand et al. 1980; Kiang 1987), which was integrated in linkage group A2 (chromosome 8) of the USDA/Iowa State University soybean molecular linkage map (Cregan et al. 1999).

β -conglycinin (7S globulin) and glycinin (11S globulin) are the major components of storage protein in soybean. β -conglycinin consists of three subunits, α' , α , β (Thanh and Shibasaki 1976) and exhibits poorer nutritional and food processing properties than glycinin. Also, β -conglycinin contains much less sulfur-containing amino acid, methionine and cysteine, than glycinin (Koshiyama 1968). Several mutant lines affecting accumulation of β -conglycinin

S. W. Choi • J. H. Park • J. I. Chung (✉)
Department of Agronomy, Gyeongsang National University,
Jinju 52828, Korea
e-mail: jongil@gnu.ac.kr

Table 1 Seed coat color, hilum color and genotype for *Cgyl* (*cgyl*) and *Ti* (*ti*) alleles of the two parents used in this experiment

Parents	Trait/genotype					
	Seed coat color	Hilum color	KTI protein	<i>Ti(ti)</i>	α' -subunit protein	<i>Cgyl</i> (<i>cgyl</i>)
BL-1	Yellow	Yellow	Absent	<i>titi</i>	Present	<i>CgylCgyl</i>
15G1	Yellow	Yellow	Present	<i>TiTl</i>	Absent	<i>cgylcgyl</i>

have been identified in soybean germplasms. Kitamura and Kaizuma (1981) identified Keburi, which was characterized by the absence of the α' -subunit of β -conglycinin. Kitamura et al. (1984) reported that the absence of α' -subunit were controlled by single recessive alleles, *cgyl*. Wild soybean line, QT2, which lacks α' -subunit was also identified. In the QT2 line, the deficiency of α' -subunit is controlled by a single dominant gene, *Scg* (Hajika et al. 1998). *Scg* gene was inherited independently with *cgyl* gene. Studies also showed that absence of the α' -subunit in QT2 was inherited independently from the presence of lipoxigenase isozymes. Hayashi et al. (2000) identified AFLP marker to be tightly linked to the gene for deficiency of the β -conglycinin. Highly negative correlation between the contents of β -conglycinin and glycinin was reported by Ogawa et al. (1989). Breeding of soybean containing large amount of glycinin compared with current varieties is possible by selecting strain of soybean that does not exhibit or lacks α' , α , and β -subunits of β -conglycinin. KTI and α' protein limit the utilization of raw soybeans as direct feed requiring a heating step to denature the activity and energy costs as well as altering the physical properties of the soybean proteins. Also excessive heat treatment may lower amino acid availability. The genetic elimination of these factors could be an alternative to the heat treatment. Development of new soybean cultivar with both KTI and α' -subunit protein free is needed to improve the nutrition values and to modify the food processing properties of soybeans. The objective of this research was to develop a new soybean strains with KTI and α' -subunit protein free (*titicgylcgyl* genotype) and proper agronomic traits.

Materials and Methods

Breeding population

Two parents were used to develop breeding population. BL-1 parent has *Cgyl* gene (α' -subunit of β -conglycinin protein present, *CgylCgyl* genotype) and *ti* gene (KTI protein absent, *titi* genotype). 15G1 parent has *cgyl* gene (α' -subunit of β -conglycinin protein absent, *cgylcgyl* genotype)

and *Ti* gene (KTI protein present, *TiTl* genotype). Seed traits and genotypes for *Cgyl* (*cgyl*) and *Ti* (*ti*) alleles of two parents are presented in Table 1. The seeds of BL-1 and 15G1 parents were planted to cross in a greenhouse. The crosses of BL-1 (*CgylCgyltiti*) x 15G1 (*cgylcgylTiTi*) were made and F₁ seeds were obtained. F₁ seeds obtained were planted in greenhouse. F₁ hybridity was confirmed on morphological traits. F₂ seeds from F₁ plant confirmed were harvested. The F₂ seeds were analyzed to screen the seed with *cgylcgyltiti* genotype (KTI and α' -subunit protein free).

Determination of α' -subunit protein by SDS-PAGE

Crude protein from parent and each F₂ seeds extracted to determine the presence or absence of α' -subunit protein of β -conglycinin electrophoretically. A piece of cotyledon from parent and each F₂ seed was removed and was incubated for 30 min (at room temperature) in 1 ml Tris-HCl, pH8.0, containing 1.56% v/v β -mercaptoethanol. After centrifugation, 50 μ l of the supernatant was added to an equivalent amount of 5X sample buffer [10% w/v sodium dodecyl sulfate (SDS), 50% v/v glycerol, 1.96% v/v β -mercaptoethanol, 1M Tris-HCl, pH 6.8]. The samples were boiled at 97°C for 5 min and then centrifuged. Two microliters of the supernatant were loaded on a 12% acrylamide SDS polyacrylamide gel electrophoresis (SDS-PAGE) medium gels in Owl Separation Systems Inc (Model:P9DS, Portsmouth, NH, USA). Electrophoresis was performed at 120 V for 7 hrs. Gels were stained overnight in an aqueous solution of 0.25 g Coomassie blue R250, 10% acetic acid, and 45% methanol. The gels were then destained with destaining solution (5% acetic acid, 14% methanol) for several hours. A Wide-Range SDS-PAGE molecular mass standard (Sigma MarkerTM, Product Code: M4038, St.Louis MO, USA) containing the 72kDa (for α' -subunit) was used to aid recognition of samples lacking the α' -subunit of β -conglycinin protein.

Determination of KTI protein by western blot analysis

Proteins of parent and each F₂ seed were separated by 10%

Fig. 1 Scheme for development of new soybean strains with *titicgy₁cgy₁* genotype (KTI and α' -subunit protein free)

or 12% SDS-PAGE, and transferred onto Immobilon-Pmembrane (PVDF, Millipore). After blocking for 2hr in TBS buffer [20mM Tris(pH7.5), 150mM NaCl, and 0.1% Tween20] with 5% nonfat dried milk (Carnation, Glendale, CA) at room temperature, the membrane were incubated with KTI antibody for 1hr. After washing in TBS buffer three times, the blot was incubated with a horseradish peroxidase conjugated secondary antibody, and the complex was visualized using an enhanced chemiluminescence kit (Amersham, Bucking-hamshire, UK). The thickness of band was then determined visually.

Selection of *titicgy₁cgy₁* genotype (KTI and α' -subunit protein free)

The F₂ seeds with *titicgy₁cgy₁* genotype (KTI and α' -subunit protein free) were planted to advance F₂ plant generation. F₂ plants with a proper agronomic traits were individually harvested. Two strains with a proper agronomical traits were advanced to F₄ plant generation. Random F₅ seeds from each F₄ plant harvested were used to confirm absence for both KTI and α' -subunit proteins. Plant height and 100-seed weight was recorded on the F₅ plant generation. Mean values of plant height and 100-seed weight were

Fig. 2 Segregation of α' -subunit protein in the parents and F₂ seeds. Arrow is the α' -subunit protein band. S: marker, P1: BL-1, P2: 15G1. +, -: presence and absence of α' -subunit protein, respectively

Fig. 3 Segregation of the Kunitz trypsin inhibitor (KTI) protein in the parents and F₂ seeds. Arrow is the KTI protein band. P1: BL-1, P2: 15G1. +, -: presence and absence of the KTI protein, respectively

compared by Duncan's multiple range test at the 5% level. Scheme for development of new soybean strain with *titicgy₁cgy₁* genotype is presented in Figure 1.

Results and Discussion

A total 168 F₂ seeds from the cross of BL-1 and 15G1 parents were obtained and analyzed for the segregation of Kunitz trypsin inhibitor (KTI) protein and α' -subunit protein of β -conglycinin. A part of the SDS-PAGE pattern for α' -subunit protein that appeared in the parents and F₂ seed generation is shown in Figure 2.

Bands for α' -subunit of β -conglycinin were segregated in F₂ seeds. Segregation pattern for Kunitz trypsin inhibitor (KTI) protein that appeared in the parents and F₂ seed generation is shown in Figure 3.

The segregation data for KTI protein and α' -subunit protein of β -conglycinin protein in the F₂ seed generation are presented in Table 2.

The segregation ratios of 9 : 3 : 3 : 1 (104 *Ti_Cgy₁* : 30

Table 2 Segregation for the presence or absence of Kunitz trypsin inhibitor (KTI) and α' -subunit proteins in F₂ seed generation

Seed protein		Seed number		χ^2 value (9:3:3:1)	P
KTI	α' -subunit	Observed	Expected		
+	+	104	94.5	5.12	0.5 - 0.1
+	-	30	31.5		
-	+	21	31.5		
-	-	13	10.5		

+, -: presence and absence of protein.

Table 3 Agronomic traits of parents and two new strains selected in this experiment

Parent/ strain	Plant height (cm)	Seed weight (g/100 seed)	Seed coat color	Hilum color
BL-1	63a ²	24.5a	Yellow	Yellow
15G1	75b	32.1b	Yellow	Yellow
S1	65a	29.2b	Yellow	Yellow
S2	66a	26.1a	Yellow	Yellow

²Different letters in the column are significantly different by DMRT at 5%.

Fig. 4 Confirmation of the α' -subunit protein free (A) and Kunitz trypsin inhibitor (KTI) protein free (B) in the parents and new strains. Arrows are the α' -subunit band (72 kDa) and KTI protein band (21.5 kDa). P1: BL-1, P2: 15G1, S1, S2: new strain (*titicgy₁cgv₁* genotype). +, -: presence and absence of α' -subunit and KTI proteins, respectively

Ti₁cgv₁cgv₁: 21 *cgv₁cgv₁Ti₁*: 13 *titicgy₁cgv₁*) between *Ti* gene and *Cgv₁* gene in F₂ seeds were observed ($\chi^2 = 5.12$, $P = 0.5-0.10$). This data showed that *Ti* gene was inherited independently with the *Cgv₁* gene in soybean. These results supported the previous reports (Kitamura et al. 1984; Orf and Hymowitz 1979; Sung et al. 2010). Only thirteen F₂ seeds with recessive allele *titicgy₁cgv₁* genotype were planted and Ten F₂ plants were individually harvested. Agronomic traits such as plant type, plant height, maturity date, lodging, seed quality, and 100-seed weight were evaluated on the ten F₃ plant strains with *titicgy₁cgv₁* genotype. Two F₄ plant strains with a proper agronomical traits were selected and harvested. Random F₅ seeds from each F₄ plant strain were used to confirm absence for both KTI and α' -subunit proteins (Fig. 4).

Kunitz trypsin inhibitor (KTI) and α' -subunit proteins were not observed in the mature F₅ seed of two new strains. Agronomic traits of two parents and two new strains with *titicgy₁cgv₁* genotype are presented in Table 3.

Two strains (S1 and S2) have yellow seed coat and hilum. Plant height of S1 strain was 65 cm compared to the parents of 63~75 cm. The 100-seed weight of S1 strain was 29.2 g compared to the parents of 24.5~32.1 g. Plant height of S2 strain was 66 cm compared to the

Fig. 5 Plants (up) and seeds (down) of the two new strains (S1 and S2) with *titicgy₁cgv₁* genotype

parents of 63~75 cm. The 100-seed weight of S1 strain was 26.1 g compared to the parents of 24.5~32.1 g. Plants and seeds of two new strains with *titicgy₁cgv₁* genotype are shown in Figure 5.

Two strains selected in this research will be used to improve new cultivar with free of KTI and α' -subunit proteins.

Reference

- Cregan PB, Jarvik T, Bush AL, Shoemaker RC, Lark KG, Kahler AL, Kaya N, VanToai TT, Lohnes DG, Chung JI, Specht JE (1999) An integrated genetic linkage map of the soybean. *Crop Sci* 39:1464-1490
- Hajika M, Takahashi M, Sakai S, Matsunaga R (1998) Dominant inheritance of a trait lacking β -conglycinin detected in a wild soybean line. *Breeding Science* 48:383-386
- Hayashi M, Nishioka M, Kitamura K, Harada K (2000) Identification of AFLP markers tightly linked to the gene for deficiency of the 7S globulin in soybean seed and characterization of abnormal phenotypes involved in the mutation.

- Breeding Science 50:123-129
- Hildebrand DF, Orf JH, Hymowitz T (1980) Inheritance of an acid phosphatase and its linkage with the Kunitz trypsin inhibitor seed protein of soybeans. *Crop Sci* 20:83-85
- Hymowitz T, Hadley HH (1972) Inheritance of a trypsin variant in seed protein of soybeans. *Crop Sci* 12:197-198
- Kiang YT (1987) Mapping three protein loci on a soybean chromosome. *Crop Sci* 27:44-46
- Kitamura K, Kaizuma N (1981) Mutant strains with low level of subunits of 7S globulins in soybean seed. *Japan J Breed* 31:353-359
- Kitamura K, Davies CS, Nielsen NC (1984) Inheritance of alleles for *Cgy₁* and *Gy₄* storage protein genes in soybean. *Theor Appl Genet* 68:253-257
- Koshiyama I (1968) Chemical and physical properties in soybean globulins. *Cereal Chem* 45:394-404
- Kunitz M (1945) Crystallization of a trypsin inhibitor from soybean. *Science* 101:668-669
- Ogawa TE, Tayama E, Kitamura K, Kaizuma N (1989) Genetic improvement of seed storage proteins using three variant alleles of 7S globulin subunits in soybean. *Japan J Breed* 39:137-147
- Orf JH, Hymowitz T (1979) Inheritance of the absence of the trypsin inhibitor in seed protein of soybeans. *Crop Sci* 19:107-109
- Singh L, Wilson CM, Hadley HH (1969) Genetic differences in soybean trypsin inhibitors separated by disc electrophoresis. *Crop Sci* 9:489-491
- Sung MK, Han EH, Kim KR, Park JS, Hwang KJ, Nam JW, Chung JI (2010) Inheritance of *Cgy₁* gene and *Ti* gene in Mature Soybean Seed. *Korean J Breed Sci* 42(1):35-39
- Thanh VH, Shibasaki K (1976) Heterogeneity of beta-conglycinin. *Biochim Biophys Acta* 469:326-328